

PALM Intranet

Application
Number **Submit**

IDS Flag Clearance for Application 10613975

IDS
Information

Content	Mailroom Date	Entry Number	IDS Review	Last Modified	Reviewer
M844	2004-02-02	15	Y <input checked="" type="checkbox"/>	2004-03-16 13:59:24.0	lhumes
Update					



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Reg]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

B

Search for

Limits

Preview/Index

History

Clipboard

Details

Display

Show

Sort by

Send to

About Entrez

Text Version

All: 31

Review: 0

Items 21 - 31 of 31

Previous of 2

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

- ☐ 21: [Bondre S, Lewandrowski KU, Hasirci V, Cattaneo MV, Gresser JD, Wise DL, Tomford WW, Trantolo DJ.](#) Related Articles, Links



Biodegradable foam coating of cortical allografts.

Tissue Eng. 2000 Jun;6(3):217-27.

PMID: 10941216 [PubMed - indexed for MEDLINE]

- ☐ 22: [Bryan DJ, Holway AH, Wang KK, Silva AE, Trantolo DJ, Wise D, Summerhayes IC.](#) Related Articles, Links



Influence of glial growth factor and Schwann cells in a bioresorbable guidance channel on peripheral nerve regeneration.

Tissue Eng. 2000 Apr;6(2):129-38.

PMID: 10941208 [PubMed - indexed for MEDLINE]

- ☐ 23: [Lewandrowski KU, Gresser JD, Wise DL, Trantolo DJ, Hasirci V.](#) Related Articles, Links



Tissue responses to molecularly reinforced polylactide-co-glycolide implants.

J Biomater Sci Polym Ed. 2000;11(4):401-14.

PMID: 10903038 [PubMed - indexed for MEDLINE]

- ☐ 24: [Lewandrowski KU, Bondre S, Gresser JD, Silva AE, Wise DL, Trantolo DJ.](#) Related Articles, Links



Augmentation of osteoinduction with a biodegradable poly(propylene glycol-co-fumaric acid) bone graft extender. A histologic and histomorphometric study in rats.

Biomed Mater Eng. 1999;9(5-6):325-34.

PMID: 10822488 [PubMed - indexed for MEDLINE]

- ☐ 25: [Lewandrowski KU, Bondre SP, Gresser JD, Wise DL, Tomford WW, Trantolo DJ.](#) Related Articles, Links



Improved osteoconduction of cortical bone grafts by biodegradable foam coating.

Biomed Mater Eng. 1999;9(5-6):265-75.

PMID: 10822482 [PubMed - indexed for MEDLINE]

- ☐ 26: [Lewandrowski KU, Gresser JD, Wise DL, White RL, Trantolo DJ.](#) Related Articles, Links



Osteoconductivity of an injectable and bioresorbable poly(propylene glycol-co-fumaric acid) bone cement.

Biomaterials. 2000 Feb;21(3):293-8.

PMID: 10646946 [PubMed - indexed for MEDLINE]

- ☐ 27: [Lewandrowski KU, Cattaneo MV, Gresser JD, Wise DL, White RL, Bonassar L, Trantolo DJ.](#) Related Articles, Links
Effect of a poly(propylene fumarate) foaming cement on the healing of bone defects.
Tissue Eng. 1999 Aug;5(4):305-16.
PMID: 10477853 [PubMed - indexed for MEDLINE]
- ☐ 28: [Hsu YY, Gresser JD, Trantolo DJ, Lyons CM, Gangadharam PR, Wise DL.](#) Related Articles, Links
Effect of polymer foam morphology and density on kinetics of in vitro controlled release of isoniazid from compressed foam matrices.
J Biomed Mater Res. 1997 Apr;35(1):107-16.
PMID: 9104703 [PubMed - indexed for MEDLINE]
- ☐ 29: [Hsu YY, Gresser JD, Stewart RR, Trantolo DJ, Lyons CM, Simons GA, Gangadharam PR, Wise DL.](#) Related Articles, Links
Mechanisms of isoniazid release from poly(d,l-lactide-co-glycolide) matrices prepared by dry-mixing and low density polymeric foam methods.
J Pharm Sci. 1996 Jul;85(7):706-13.
PMID: 8818994 [PubMed - indexed for MEDLINE]
- ☐ 30: [Gresser JD, Hsu SH, Nagaoka H, Lyons CM, Nieratko DP, Wise DL, Barabino GA, Trantolo DJ.](#) Related Articles, Links
Analysis of a vinyl pyrrolidone/poly(propylene fumarate) resorbable bone cement.
J Biomed Mater Res. 1995 Oct;29(10):1241-7.
PMID: 8557726 [PubMed - indexed for MEDLINE]
- ☐ 31: [Jimoh AG, Wise DL, Gresser JD, Foote RH, Rhodes RC, Underbill LH, Trantolo DJ.](#) Related Articles, Links
Pulsatile release of FSH for superovulation in cattle.
Theriogenology. 1995 Feb;43(3):645-56.
PMID: 16727656 [PubMed - in process]

Items 21 - 31 of 31

Previous **Page** 2 of 2Display **Summary** Show **20** Sort by Send to[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 30 2007 04:56:27

A service of the National Library of Medicine
and the National Institutes of HealthMy NCBI
[Sign In] [Reg]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

B

Search PubMed for wise and trantolo

Limits

Preview/Index

History

Clipboard

Details

Display

Summary

Show

20

Sort by

Send to

About Entrez

Text Version

All: 31

Review: 0

Items 1 - 20 of 31

Page

1

of 2 Next

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

- ☐ 1: [Bryan DJ, Tang JB, Doherty SA, Hile DD, Trantolo DJ, Wise DL, Summerhayes IC.](#) Related Articles, Links



Enhanced peripheral nerve regeneration through a poled bioresorbable poly (lactic-co-glycolic acid) guidance channel.

J Neural Eng. 2004 Jun;1(2):91-8. Epub 2004 Jun 14.

PMID: 15876627 [PubMed - indexed for MEDLINE]

- ☐ 2: [Peacock ZS, Barnes LA, King WF, Trantolo DJ, Wise DL, Taubman MA, Smith DJ.](#) Related Articles, Links



Influence of microparticle formulation on immunogenicity of SYI, a synthetic peptide derived from Streptococcus mutans GbpB.

Oral Microbiol Immunol. 2005 Feb;20(1):60-4.

PMID: 15612949 [PubMed - indexed for MEDLINE]

- ☐ 3: [Hasirci V, Litman AE, Trantolo DJ, Gresser JD, Wise DL, Margolis HC.](#) Related Articles, Links



PLGA bone plates reinforced with crosslinked PPF.

J Mater Sci Mater Med. 2002 Feb;13(2):159-67.

PMID: 15348638 [PubMed]

- ☐ 4: [Smith DJ, Lam A, Barnes LA, King WF, Peacock Z, Wise DL, Trantolo DJ, Taubman MA.](#) Related Articles, Links



Remote glucosyltransferase-microparticle vaccine delivery induces protective immunity in the oral cavity.

Oral Microbiol Immunol. 2003 Aug;18(4):240-8.

PMID: 12823800 [PubMed - indexed for MEDLINE]

- ☐ 5: [Hile DD, Kirker-Head C, Doherty SA, Kowaleski MP, McCool J, Wise DL, Trantolo DJ.](#) Related Articles, Links



Mechanical evaluation of a porous bone graft substitute based on poly (propylene glycol-co-fumaric acid).

J Biomed Mater Res B Appl Biomater. 2003 Jul 15;66(1):311-7.

PMID: 12808589 [PubMed - indexed for MEDLINE]

- ☐ 6: [Lewandrowski KU, Bondre SP, Wise DL, Trantolo DJ.](#) Related Articles, Links




Enhanced bioactivity of a poly(propylene fumarate) bone graft substitute by augmentation with nano-hydroxyapatite.

Biomed Mater Eng. 2003;13(2):115-24.

PMID: 12775902 [PubMed - indexed for MEDLINE]

- ☐ 7: [Trantolo DJ, Sonis ST, Thompson BM, Wise DL, Lewandrowski](#) Related Articles, Links


KU, Hile DD.

-  Evaluation of a porous, biodegradable biopolymer scaffold for mandibular reconstruction.

Int J Oral Maxillofac Implants. 2003 Mar-Apr;18(2):182-8.

PMID: 12705295 [PubMed - indexed for MEDLINE]


- ☐ 8: Lewandrowski KU, Bondre SP, Shea M, Untch CM, Hayes WC, Hile DD, Wise DL, Trantolo DJ. Related Articles, Links

-  Composite resorbable polymer/hydroxylapatite composite screws for fixation of osteochondral osteotomies.

Biomed Mater Eng. 2002;12(4):423-38.

PMID: 12652036 [PubMed - indexed for MEDLINE]


- ☐ 9: Bryan DJ, Tang JB, Holway AH, Rieger-Christ KM, Trantolo DJ, Wise DL, Summerhayes IC. Related Articles, Links

-  Enhanced peripheral nerve regeneration elicited by cell-mediated events delivered via a bioresorbable PLGA guide.

J Reconstr Microsurg. 2003 Feb;19(2):125-34.

PMID: 12632314 [PubMed - indexed for MEDLINE]


- ☐ 10: Lewandrowski KU, Hile DD, Thompson BM, Wise DL, Tomford WW, Trantolo DJ. Related Articles, Links

-  Quantitative measures of osteoinductivity of a porous poly(propylene fumarate) bone graft extender.

Tissue Eng. 2003 Feb;9(1):85-93.

PMID: 12625957 [PubMed - indexed for MEDLINE]


- ☐ 11: Lewandrowski KU, Bondre S, Hile DD, Thompson BM, Wise DL, Tomford WW, Trantolo DJ. Related Articles, Links

-  Porous poly(propylene fumarate) foam coating of orthotopic cortical bone grafts for improved osteoconduction.

Tissue Eng. 2002 Dec;8(6):1017-27.

PMID: 12542947 [PubMed - indexed for MEDLINE]


- ☐ 12: Lewandrowski KU, Bondre SP, Shea M, Untch CM, Hayes WC, Hile DD, Wise DL, Trantolo DJ. Related Articles, Links

-  Composite poly(lactide)/hydroxylapatite screws for fixation of osteochondral osteotomies. A morphometric, histologic and radiographic study in sheep.

J Biomater Sci Polym Ed. 2002;13(11):1241-58.

PMID: 12518802 [PubMed - indexed for MEDLINE]


- ☐ 13: Lewandrowski KU, Bondre SP, Wise DL, Trantolo DJ. Related Articles, Links

-  Healing of osteochondral osteotomies after fixation with a hydroxyapatite-buffered polylactide. A histomorphometric and radiographic study in rabbits.

Biomed Mater Eng. 2002;12(3):259-70.

PMID: 12446941 [PubMed - indexed for MEDLINE]







- ☐ 14: Kandziora F, Pflugmacher R, Kleemann R, Duda G, Wise DL, Trantolo DJ, Lewandrowski KU. Related Articles, Links

-  Biomechanical analysis of biodegradable interbody fusion cages augmented With poly(propylene glycol-co-fumaric acid).

Spine. 2002 Aug 1;27(15):1644-51.

PMID: 12163726 [PubMed - indexed for MEDLINE]

Hasirci V, Berthiaume F, Bondre SP, Gresser JD, Trantolo DJ,

- ☐ 15: [Toner M, Wise DL](#). Related Articles, Links
 Expression of liver-specific functions by rat hepatocytes seeded in treated poly(lactic-co-glycolic) acid biodegradable foams.
Tissue Eng. 2001 Aug;7(4):385-94.
PMID: 11506728 [PubMed - indexed for MEDLINE]
- ☐ 16: [Smith DJ, King WF, Barnes LA, Trantolo D, Wise DL, Taubman MA](#). Related Articles, Links
 Facilitated intranasal induction of mucosal and systemic immunity to mutans streptococcal glucosyltransferase peptide vaccines.
Infect Immun. 2001 Aug;69(8):4767-73.
PMID: 11447149 [PubMed - indexed for MEDLINE]
- ☐ 17: [Hasirci V, Lewandrowski K, Gresser JD, Wise DL, Trantolo DJ](#). Related Articles, Links
 Versatility of biodegradable biopolymers: degradability and an in vivo application.
J Biotechnol. 2001 Mar 30;86(2):135-50.
PMID: 11245902 [PubMed - indexed for MEDLINE]
- ☐ 18: [Lewandrowski KU, Gresser JD, Bondre S, Silva AE, Wise DL, Trantolo DJ](#). Related Articles, Links
 Developing porosity of poly(propylene glycol-co-fumaric acid) bone graft substitutes and the effect on osteointegration: a preliminary histology study in rats.
J Biomater Sci Polym Ed. 2000;11(8):879-89.
PMID: 11211098 [PubMed - indexed for MEDLINE]
- ☐ 19: [Smith DJ, Trantolo DJ, King WF, Gusek EJ, Fackler PH, Gresser JD, De Souza VL, Wise DL](#). Related Articles, Links
 Induction of secretory immunity with bioadhesive poly (D,L-lactide-co-glycolide) microparticles containing Streptococcus sobrinus glucosyltransferase.
Oral Microbiol Immunol. 2000 Apr;15(2):124-30.
PMID: 11155176 [PubMed - indexed for MEDLINE]
- ☐ 20: [Hasirci V, Lewandrowski KU, Bondre SP, Gresser JD, Trantolo DJ, Wise DL](#). Related Articles, Links
 High strength bioresorbable bone plates: preparation, mechanical properties and in vitro analysis.
Biomed Mater Eng. 2000;10(1):19-29.
PMID: 10950204 [PubMed - indexed for MEDLINE]

Items 1 - 20 of 31

Page

1

of 2 Next

Display

Summary



Show

20



Sort by



Send to



[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 30 2007 04:56:27



US Patent and Trademark Office -
Washington

User name:

Password:

[Register](#) [Forgotten Password](#) [Athens Login](#)

[Synergy Home](#) | [Browse](#) | [Search](#) | [My Synergy](#) | [Books Online](#) | [Resources](#) |
[About](#) | [Help](#)

ORAL MICROBIOLOGY AND IMMUNOLOGY

Journal Menu

[Table of Contents](#)
[List of Issues](#)

Tools

[Email this article](#)
[Add to favorite articles](#)
[Export this citation](#)
[Alert me when this article is
cited: Email | RSS](#)
(What is this?)

[View PubMed citation](#)
[View ISI citation](#)

Publication history

Issue online:
25 Dec 2001

Accepted for publication April
26, 1999

[Home](#) > [List of Issues](#) > [Table of Contents](#) > Article Abstract

Oral Microbiology and Immunology


Volume 15 Issue 2 Page 124 - April 2000

To cite this article: D. J. Smith, D. J. Trantolo, W.
F. King, E. J. Gusek, P. H. Fackler, J. D. Gresser, V.
L. De Souza, D. L. Wise (2000)

Induction of secretory immunity with bioadhesive poly (D,L-lactide-
co-glycolide) microparticles containing *Streptococcus sobrinus*
glucosyltransferase

Oral Microbiology and Immunology 15 (2), 124-130.

doi:10.1034/j.1399-302x.2000.150209.x



[Prev Article](#) | [Next Article](#)

Abstract

Induction of secretory immunity with bioadhesive poly (D,L-lactide-co-glycolide) microparticles containing *Streptococcus sobrinus* glucosyltransferase

D. J. Smith¹, D. J. Trantolo², W. F. King¹, E. J. Gusek², P. H. Fackler²,
J. D. Gresser², V. L. De Souza¹, D. L. Wise²

¹Department of Immunology, Forsyth Dental Center, Boston,

²Cambridge Scientific, Inc., Belmont, Massachusetts, USA

Daniel J. Smith, Forsyth Dental Center, 140 The Fenway, Boston,
MA 02115, USA

Abstract

The effect of mucosal delivery of *Streptococcus sobrinus* glucosyltransferase (GTF) in bioadhesive poly (D,L-lactide-co-glycolide) (PLGA) microparticles on induction of salivary IgA and serum IgG antibody responses was measured in Sprague-Dawley rats. Preparations of GTF/PLGA/gelatin microparticles, or PLGA/gelatin microparticles or GTF in alum, were administered four times at weekly intervals by intranasal or intragastric routes. Two subcutaneous injections of GTF in PLGA/gelatin microparticles or in alum were given to separate groups of rats. Significant elevations in salivary IgA antibody levels to *S. sobrinus* GTF were observed only in the groups

immunized intranasally 28 days after immunizations were begun. Five of six rats given the GTF microparticles intranasally had positive salivary IgA antibody responses to GTF, and the mean salivary IgA antibody level of this group was 30-fold higher than any other mucosally or systemically immunized group. Salivary IgA responses in the GTF-microparticle group remained significantly higher than all other mucosally immunized groups for at least 10 weeks after the primary immunization. All rats in this group demonstrated aspects of anamnesis following a more limited secondary course of intranasal administration. Intranasal administration of GTF in microparticles also induced a serum IgG response to GTF in some rats. After secondary intranasal GTF microparticle administration, several rats had sustained serum IgG antibody levels that were within the range of sera from rats subcutaneously injected with GTF in microparticles or in alum. Thus intranasal delivery of GTF-containing bioadhesive microparticles induced the highest and longest lasting salivary immune response of any mucosal or systemic route or vehicle tested and could be expected to be a useful method for induction of mucosal immunity.

This article is cited by:

- Z. S. Peacock, L. A. Barnes, W. F. King, D. J. Trantolo, D. L. Wise, M. A. Taubman, D. J. Smith. (2005) Influence of microparticle formulation on immunogenicity of SYI, a synthetic peptide derived from *Streptococcus mutans* GbpB. *Oral Microbiology and Immunology* **20**:1, 60–64
[Abstract](#) [Abstract and References](#) [Full Text Article](#) [Full Article PDF](#)
- D. J. Smith, A. Lam, L. A. Barnes, W. F. King, Z. Peacock, D. L. Wise, D. J. Trantolo, M. A. Taubman. (2003) Remote glucosyltransferase-microparticle vaccine delivery Induces protective Immunity in the oral cavity. *Oral Microbiology and Immunology* **18**:4, 240–248
[Abstract](#) [Abstract and References](#) [Full Text Article](#) [Full Article PDF](#)
- D.J. Aframian, R.S. Redman, S. Yamano, J. Nikolovski, E. Cukierman, K.M. Yamada, M.F. Kriete, W.D. Swaim, D.J. Mooney, B.J. Baum. (2002) Tissue Compatibility of Two Biodegradable Tubular Scaffolds Implanted Adjacent to Skin or Buccal Mucosa in Mice. *Tissue Engineering* **8**:4, 649
[CrossRef](#)

This Article

Abstract

References

Full Text HTML

Full Text PDF (94 KB)

Rights & Permissions

Search

In

- ☒ Synergy
- ☐ PubMed (MEDLINE)
- ☐ CrossRef

By keywords

- ☐ PLGA
- ☐ microparticles
- ☐ glucosyltransferase
- ☐ *Streptococcus sobrinus*
- ☐ saliva

☐ IgA

By author

☐ D. J. Smith

☐ D. J. Trantolo

☐ W. F. King

☐ E. J. Gusek

☐ P. H. Fackler

☐ J. D. Gresser

☐ V. L. De Souza

☐ D. L. Wise

☐

GO 

[Privacy Statement](#) | [Terms & Conditions](#) | [Contact](#) | [Help](#)



Blackwell Synergy® is a Blackwell Publishing, Inc. registered trademark Technology Partner — [Atypon Systems, Inc.](#)
Partner of CrossRef, COUNTER, AGORA, HINARI and OARE



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Regis]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

☐ 1: Tissue Eng. 2000 Jun;6(3):217-27.

Mary Ann Liebert

Links

Biodegradable foam coating of cortical allografts.

Bondre S, Lewandrowski KU, Hasirci V, Cattaneo MV, Gresser JD, Wise DL, Tomford WW, Trantolo DJ.

Department of Chemical Engineering, Northeastern University, Boston, Massachusetts, USA.

Clinical outcomes of bone allograft procedures may be improved by modifying the surface of the graft with an osteoconductive biopolymeric coating. In this comparative in vitro study, we evaluated the dimensional stability, mechanical strength, hydrophilicity, and water uptake of biodegradable foams of poly(propylene fumarate) (PPF) and poly(D,L-lactic-co glycolic acid) (PLGA) when applied as surface coatings to cortical bone. Cortical bone samples were divided into four groups: Type I, untreated bone; Type II, laser-perforated bone; Type III, partially demineralized bone; and Type IV, laser-perforated and partially demineralized bone. Results show that PPF wets easily, achieving 12.5% wt/wt in 30 min. Compressive tests on the PPF foam material showed that the compressive strength was 6.8 MPa prior to in vitro incubation but then gradually reduced to 1.9 MPa at 8 weeks. Push-out and pulloff strength tests showed that initially both PPF and PLGA foam coatings had comparable adherence strengths to the cortical bone samples (100-150 N). When additional geometrical surface alteration by perforation and demineralization of the bony substrate was employed, in vitro adherence of the PPF foam coating was further increased to 120 N, demonstrating a statistically significant improvement of push-out strength throughout the entire 8-week observation period ($p < 0.0002$ for all four data points). The pore geometry of PPF-foam coatings changed little over the 2-month evaluation period. In comparison, PLGA foam coating around the cortical bone samples rapidly lost structure with a decrease of 67% in strength seen after 1-week in vitro incubation. These new types of bone allografts may be particularly useful where the use of other replacement materials is not feasible or practical.

PMID: 10941216 [PubMed - indexed for MEDLINE]

Related Links

Porous poly(propylene fumarate) foam coating of orthotopic cortical bone grafts for improved osteoconduction. [Tissue Eng. 2002]

Improved osteoconduction of cortical bone grafts by biodegradable foam coating. [Biomed Mater Eng. 1999]

Controlled drug release from a novel injectable biodegradable microsphere/scaffold composite based on poly(D,L-lactide-co-glycolide). [J Biomed Mater Res. 2000]

Versatility of biodegradable biopolymers: degradability and an in vivo application. [J Biotechnol. 2001]

Fabrication of biodegradable polymer scaffolds to engineer trabecular bone. [J Biomater Sci Polym Ed. 1995]

See all Related Articles...

Display AbstractPlus Show 20 Sort by Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 30 2007 04:56:27



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

☐ 1: J Biomater Sci Polym Ed. 2000;11(4):401-14.



Links

Tissue responses to molecularly reinforced polylactide-co-glycolide implants.

Lewandrowski KU, Gresser JD, Wise DL, Trantolo DJ, Hasirci V.

Orthopaedic Research Laboratories, Massachusetts General Hospital, Boston 02114, USA.

Plates for internal fixation fabricated from biodegradable polymers degrade via an autocatalytic route. When they are used in bone implants of significant size and thickness, hollowing of the implant may occur while the overall dimensions appear unchanged. We hypothesized that incorporation of a cross-linked polypropylene fumarate matrix into polylactide-co-glycolide bone plates may provide an internal molecular network which prevents implant collapse. Cross-linking reagents of varying hydrophilicity including N-vinylpyrrolidone (VP), hydroxyethylmethacrylate (HEMA), and ethyleneglycol dimethacrylate (EGDMA) were employed. With the objective of determining the most biocompatible and structurally sound composition for molecular reinforcement, we investigated tissue responses in both subcutaneous and orthotopic rodent implantation models in relation to maintenance of implant integrity by histologic, histomorphometric, and stereomicroscopic analysis. Results showed that tissue responses were correlated with dimensional stability of the implants. The most favorable results were seen with the hydrophobic cross-linker EGDMA; this may have been related to the initial reduction of the water uptake by the implant. Cross-linking of polypropylene fumarate with EGDMA within a polylactide-co-glycolide bone plate may offer a means to maintain excellent biocompatibility while improving dimensional stability of biodegradable bone plates.

PMID: 10903038 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Related Links

High strength bioresorbable bone plates: preparation, mechanical properties and [Biomater Mater Res 2000](#)

Versatility of biodegradable biopolymers: degradability and an in vivo application. [\[J Biotechnol. 2001\]](#)

Comparison of the pull-out forces of bioabsorbable polylactide/glycolide screws (Biosorb and Lactosorb) and tacks: a study on the stability of fixation in human cadaver parietal bones. [\[J Craniofac Surg. 2002\]](#)

In vivo evaluation of a novel electrically conductive polypyrrole/poly(D,L-lactide) composite and polypyrrole-coated poly(D,L-lactide-co-glycolide) membrane [\[J Biomed Mater Res A. 2004\]](#)

PLGA bone plates reinforced with crosslinked [\[J Biomater Sci Mater Med. 2002\]](#)

[See all Related Articles...](#)

Write to the Help Desk
NCBI | NLM | NIH

Department of Health & Human Services



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals B

Search PubMed ☒ for wise and (DNA and foam)

Limits Preview/Index History Clipboard Details

Display AbstractPlus ☒ Show 20 ☒ Sort by ☒ Send to ☒

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Tissue Eng. 2001 Aug;7(4):385-94.

Mary Ann Liebert, Link

Expression of liver-specific functions by rat hepatocytes seeded in treated poly(lactic-co-glycolic) acid biodegradable foams.

Hasirci V, Berthiaume F, Bondre SP, Gresser JD, Trantolo DJ, Toner M, Wise DL.

Department of Biological Sciences, Biotechnology Research Center, Middle East Technical University, Ankara, Turkey.

Techniques of liver replacement would benefit patients awaiting donor livers and may be a substitute for transplantation in patients whose livers can regenerate. Poly(lactic-co-glycolic acid) (PLGA) copolymers are biodegradable and have been shown to be useful as scaffolds for seeding and culturing various types of cells. In this study, foam disks were prepared from PLGA (lactic-to-glycolic mole ratio of 85:15) by lyophilization of benzene (5% w/v) solutions. These disks were then used as scaffolds for rat hepatocyte culture. Foams were coated with either a type I collagen gel (0.1% w/v), coated with gelatin (5% w/v), or treated with oxygen plasma (25 W, 90 s) to modify their surface chemistry and wettability. The disks were then seeded with rat hepatocytes (10(6)/mL) and cultured for a period of 2 weeks. All surface treatments resulted in increased hydrophilicity, the greatest being obtained by collagen treatment (contact angle < 10 degrees), and a minimal decrease in void fraction (5%). DNA content after a 2-week culture period increased proportionally with the wettability of the treated foam surface. Urea synthesis in untreated foams averaged 15.3 +/- 2.3 microg/h/microg DNA, which was significantly higher than that for controls, whereas gelatin and collagen treated foams exhibited urea synthetic rates below the control levels at all times. The DNA content decreased significantly by about 50% between days 1 and 12. PLGA foams, treated and untreated, represent a promising scaffold for scaling up hepatocyte cultures.

Related Links

Growth and metabolism of human hepatocytes on biomodified collagen poly(lactic-co-glycolic acid) three-dimensional scaffolds. [ASAIO J. 2006]

Three-dimensional culture of rat calvarial osteoblasts in porous biodegradable polymers. [Biomaterials. 1998]

Wetting of poly(L-lactic acid) and poly(DL-lactic-co-glycolic acid) foams for tissue culture. [Biomaterials. 1994]

Adhesion behaviours of hepatocytes cultured onto biodegradable polymer surface modified by alkali hydrolysis. [Biomaterials. 1999]

In vitro and in vivo degradation of porous poly(DL-lactic-co-glycolic acid) foams. [Biomaterials. 2000]

See all Related Articles...

PMID: 11506728 [PubMed - indexed for
MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 30 2007 04:56:27